

University of Groningen

## Annual chronotypes functionally link life histories and life cycles in birds

Karagicheva, Julia; Rakhimberdiev, Eldar; Saveliev, Anatoly; Piersma, Theunis

*Published in:*  
Functional Ecology

*DOI:*  
[10.1111/1365-2435.13181](https://doi.org/10.1111/1365-2435.13181)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Final author's version (accepted by publisher, after peer review)

*Publication date:*  
2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Karagicheva, J., Rakhimberdiev, E., Saveliev, A., & Piersma, T. (2018). Annual chronotypes functionally link life histories and life cycles in birds. *Functional Ecology*, 32(10), 2369-2379.  
<https://doi.org/10.1111/1365-2435.13181>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Functional Ecology

DR JULIA KARAGICHEVA (Orcid ID : 0000-0002-2404-6826)

DR ELDAR RAKHIMBERDIEV (Orcid ID : 0000-0001-6357-6187)

Article type : Research Article

Corresponding author mail id : [jkaraj@gmail.com](mailto:jkaraj@gmail.com)

Editor: Brett Sandercock

Section: Animal Physiological Ecology

Revised manuscript for *Functional Ecology* (dd. 06 June 2018):

Annual chronotypes functionally link life histories and life cycles in birds

Julia Karagicheva<sup>1</sup>, Eldar Rakhimberdiev<sup>1,2</sup>, Anatoly Saveliev<sup>3</sup> & Theunis Piersma<sup>1,4</sup>

<sup>1</sup>*NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems and Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands*

<sup>2</sup>*Department of Vertebrate Zoology, Biological Faculty, Lomonosov Moscow State University, 119991, Moscow, Russian Federation*

<sup>3</sup>*Institute of Ecology and Geography, Kazan Federal University, Kazan, Russian Federation*

<sup>4</sup>*Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands*

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1365-2435.13181

This article is protected by copyright. All rights reserved.

## Abstract

1. Life-history responses to ecological selection pressures can be described by a slow-fast life-history axis. Along this axis, fast-living animals usually invest in high breeding output, while slow-living ones prioritize their own survival.
2. Birds may solve the trade-off between reproduction and survival by optimising their seasonal schedules. Breeding early tends to facilitate reproductive success, whereas breeding late increases the chances to survive. On the basis of this argument, short- and long-lived birds should benefit from initiating spring activities earlier and later, respectively.
3. The timing of seasonal activities, all else being equal, depends on the architecture of endogenous circannual clocks. Particularly, the length of the circannual period relative to the 365-day environmental year facilitates either the anticipation of seasonal activities (in case of periods shorter than 365 days) or represents a responsive mode (when periods are longer than 365 days). The two alternatives will be manifested by early or late annual chronotypes, respectively.
4. We hypothesise that, in birds, annual chronotype will correspond with position on the 'pace-of-life scale'. Species with low survival probability, and thus a poor chance of breeding in a next season, should show early annual chronotypes facilitated by circannual clock periods shorter than 365 days. In contrast, species with high survival rates should benefit from relatively long circannual periods.
5. We predicted that circannual period lengths should correlate positively with species-specific adult annual survival rates. Using published data for 16 wild bird species, we confirmed the predicted correlation. In our analysis, we accounted for the possible metabolic nature of circannual clocks, a correlation between rate of metabolism and survival, and phylogenetic relationships.

6. Based on our finding, we propose that evolutionary responsive circannual clocks help birds cope with temporal variation in environment in ways that are most appropriate for their life-history and life-table attributes.

Key words: annual chronotype, circannual rhythm, life-cycle stage, metabolic rate, pace of life, reproduction versus survival trade-off, seasonal migration, slow-fast life-history continuum

## Introduction

Time runs at different paces for fast- and slow-living organisms (Helm & Shavit, 2017). The ‘pace-of-life’ of organisms is reflected in the position on a virtual slow-fast life-history continuum (Ricklefs & Wikelski, 2002; Versteegh, Schwabl, Jaquier, & Tieleman, 2012).

Along the continuum, fast-living organisms tend to be smaller, mature faster, reproduce at a higher rate, and have shorter lifespans than slow-living ones. The continuum is the result of correlated variations in organismal rate processes along multiple physiological and

behavioural axes, which include growth, metabolism, sexual maturation, learning and competitive risk-taking (Bielby *et al.*, 2007; Reale *et al.*, 2010; Ricklefs & Wikelski, 2002).

When the rate processes are not intercorrelated, they are often complementary or compensatory to each other (Glazier, 2015). Such adaptive coordination facilitates an optimal match between internal and environmental conditions (Glazier, 2015; Ricklefs & Wikelski, 2002).

The adaptively coordinated rate processes in an organism form its unique life-history strategy of allocating resources among competing fitness components, especially reproduction and survival (McGraw & Caswell, 1996; McNamara, Welham, Houston, Daan, & Tinbergen, 2004; Orzack & Tuljapurkar, 2001; Sæther & Bakke, 2000). In birds, and

perhaps especially in seasonally migrating populations, the reproduction-survival trade-off is often mediated by time of a season: early breeding usually facilitates high reproductive outcome, but entails higher risks of death due to challenging weather, possible starvation due to insufficient body stores, or the risk of predation (see Drent, Both, Green, Madsen, & Piersma, 2003; Møller, 1994 for empirical examples, and Harts, Kristensen, & Kokko, 2016; Kokko, 1999; Kristensen, Johansson, Ripa, & Jonzén, 2015 for theoretical background). In contrast, breeding late is safer in terms of survival, but often decreases reproductive output (Lok, Veldhoen, Overdijk, Tinbergen, & Piersma, 2017; Weiser *et al.*, 2017). Thus, a relatively early induction of spring activities should be beneficial for species with “fast” life-histories, for which fitness depends on instantaneous reproductive success rather than a long life. The opposite should hold for birds with “slow” life histories. Therefore, when survival-reproduction trade-off is mediated by timing of breeding, resource allocation to either fitness component can be optimised by fine-tuning the timing of spring activities, as it is often found in birds breeding in seasonal environments at temperate and northern latitudes (Drent & Daan, 1980; Helm & Lincoln, 2017; Karagicheva, Liebers, *et al.*, 2016; Öberg, Pärt, Arlt, Laugen, & Low, 2014).

Spring activities, such as pre-flight fuel storage, pre-alternate moult of plumage, migratory restlessness, and the actual migratory flight, are triggered by environmental cues or *Zeitgebers* such as photoperiod, phenology or social interactions (van Wijk, Schaub, & Bauer, 2017; Williams, 2012; Bradshaw & Holzapfel, 2007; Dawson, 2008, 2015; Helm, Piersma, & van der Jeugd, 2006). Nevertheless, circannual clocks also play a crucial role in their timing (Gwinner 1986). Circannual clocks are endogenous oscillator systems suggested to be driven either by whole organism metabolism (Wikelski *et al.*, 2008) or by cyclic gonadal activity (Dawson, King, Bentley, & Ball, 2001). The clocks have a period of about 12 months, but usually run ahead or behind; they need systematic resetting by external time

cues (Gwinner 1986). In the absence of relevant cues, such as under experimental conditions, circannual clocks are not reset and run free with a wide range of periods (to our knowledge, from 6 to 18 months; see Gwinner 1986). Under constant photoperiod and temperature, some seasonal activities are exerted spontaneously, at approximately the same phase of circannual cycle as when entrained to environmental cues (reviewed in Gwinner, 1986; Helm, Schwabl, & Gwinner, 2009).

Whether exposure to environmental cues will facilitate or delay exertion of seasonal activities largely depends on the period-length relationship between the circannual oscillator and the environmental cycle (in this case: year). According to the phase-period rule, deviation of intrinsic periods from environmental cycles, i.e. the cycles of environmental *Zeitgebers*, will affect the rate of entrainment (cue-response) and determine whether the animal has an early or a late chronotype (Helm *et al.*, 2017; Helm & Lincoln, 2017; Roenneberg, Daan, & Merrow, 2003). A number of factors can affect the rate of entrainment, but in general, organisms with relatively short intrinsic cycles will be early and the ones with long cycles will be late (Floessner & Hut, 2017; Roenneberg *et al.*, 2003). Thus, animals with circannual periods shorter than 12-months will exert early ‘annual chronotypes’, and be prepared to initiate seasonal activities even before the environment becomes ‘ready’. In this case, external cues will prevent seasonal activities from being triggered too early (see Fig. 1 in Chandola-Saklani, Thapliyal, Negi, Diyundi, & Choudhary, 2004 for an example). With circannual periods longer than the environmental cycle, physiological changes in the organism will lag behind the seasonal development of the environment. As a consequence, external cues will have to facilitate the timely expression of physiological change.

We hypothesised that when timing of activities is shaped by long-term survival perspectives, short- and long-lived birds will increase their fitness by earlier or later responses to seasonal cues and will thus benefit from having short and long circannual

periods, respectively. We therefore predict that, in birds, annual chronotype should correlate with species-specific annual survival probability in the wild. To test this prediction, we compiled available published data on (1) the lengths of circannual period expressed under constant photoperiodic conditions indoors, and (2) species-specific adult annual survival rate. We then calculated the deviation of the circannual-period lengths from the 365-days environmental year and regressed the resulting representation of circannual chronotype on annual survival rate, taking into account the notion that metabolic rate correlates with both survival and circannual chronotype due to the possible metabolic nature of circannual clocks. A possible effect of shared ancestry was accounted for by including a phylogenetic correlation in the model. We discuss our results from an ecological perspective and suggest a theoretical explanation for the variation in circannual period length along the slow-fast life-history axis in birds. To our knowledge, this is the first comparative study linking the features of circannual clocks with life-history traits.

## Methods

We assembled published data on free-running circannual periods for birds maintained under experimentally fixed photoperiodic and temperature conditions in captivity. Our sample included 14 species of songbirds (order Passeriformes) and two shorebird species (order Charadriiformes), the red knot *Calidris canutus* and the great knot *C. tenuirostris* (available as Appendix S1 and at <https://git.io/vAuyo>). We had to limit our dataset to species for which data on both circannual periods and annual survival rates were available. The published data on circannual periods were found by browsing for the term “circannual” and its synonyms in [www.scholar.google.com](http://www.scholar.google.com), using reference lists in books, and by screening publications of authors specialized in chronobiology.

Data on circannual periods were available either as reported values for individual birds or as published diagrams of changes in the physiological parameters of different individuals over time. In the latter case, the beginning and the end dates of each period were estimated from the time axis of the graphic. If we had a choice of data collected in various photo-regimes, we selected for datasets from those experiments where photo-regime was kept close to 12:12 LD (12 hours light/12 hours dark). The circannual period lengths were estimated, for each individual, as the intervals between onset dates of a life-cycle stage (seasonal activity) in successive years. However, it is often impossible to unequivocally define the onset dates, since some measurements can fluctuate for many reasons. In such cases we had to apply rules of thumb. When estimating gonadal cycles from the graphics, we defined the initiation date as the time-point at which growing gonads reached a size of 2 mm (Gwinner 1981). For pre-migratory body mass peaks, the onset dates were defined as the time-point at which body mass started to steadily increase to reach maximal value.

The data included circannual periods for all available life-cycle stages. However, circannual period lengths often change in the course of the cue-deprived treatment (Gwinner, 1986; Karagicheva, Rakhimberdiev, *et al.*, 2016), and most of the circannual period lengths were available for the first one or two years spent under experimental conditions. We therefore limited our sample to transitory cycles during which the animals were transferred from natural to constant photoperiods (cycle 0), and the first cycle (cycle 1) when animals were exposed to a constant photoregime the whole year (Table 1).

We calculated deviations of circannual periods from the calendar year (hereafter, circannual deviation) for each available life-cycle stage by subtracting 365 days from the observed duration of a circannual cycle (Table 1). The species-specific adult annual survival rates in the wild (Appendix S2 and at <https://git.io/vAuSe>), metabolic rates (Appendix S3 and <https://git.io/vAuSZ>) and body mass (Appendix S4 and at <https://git.io/vAuSB>) were



searched for in [www.scholar.google.com](http://www.scholar.google.com) and in an on-line database (*COMADRE Animal Matrix Database*, 2015). If several values per species were available, we used their median in the analysis (Table 1). Survival rates of crossbills *Loxia curvirostra* were likely to be biased low because crossbills are irruptive with low site fidelity, and thus difficult to resight or recapture (Alonso & Arizaga, 2013).

Although time-keeping may differ between sexes (Hau *et al.*, 2017; Rani & Kumar, 2013), we were not able to account for this possibility as sex of experimental birds was unknown in six of the 16 species, while in nine species only males were used in the experiments. As there was no obvious trend in the proportion of males versus unknown sex along the annual survival axis, we opted not to include sex as a variable in our model.

The species in our study had an almost 20-fold range in body masses, from 8 g in the common chiffchaff *Phylloscopus collybita* to 150 g in the great knot (see Appendix S4). In comparative studies looking for functional links between longevity and other traits across species, it is generally recommended to account for species-specific body mass to eliminate the part of the correlation caused by the allometric relationship between most organismal processes and body mass (Speakman, 2005). However, organismal processes involved in time-keeping are energy- rather than mass-related (Riede, van der Vinne, & Hut, 2017; Speakman, 2005; Wikelski *et al.*, 2008). Since, in our dataset (Appendix S5 and at <https://git.io/vAu9I>), basal metabolic rate (BMR) correlated with body mass, we used BMR to account for the body-size effect. Following Speakman (2005), we regressed the circannual deviation and annual survival rate (log-transformed, natural logarithm) on the values of BMR (also log-transformed, natural logarithm) published for the 16 species (Table 1) and used the residuals as the response and explanatory variables, respectively, in models testing for the correlation between the two traits.

To control for phylogenetic autocorrelation in the analysis (Harvey & Pagel, 1991; Speakman, 2005), we introduced a phylogenetic variable in the models. We accounted for phylogenetic uncertainty (Huelsenbeck, 2000) by generating 10,000 trees with different topologies for all species in our dataset based on a comprehensive bird phylogeny for 9993 species (Hackett 'All Species' from Jetz, Thomas, Joy, Hartmann, & Mooers, 2012), using an online phylogeny generating tool (available at <http://birdtree.org/>). Different trait-evolution models require specific statistical approaches, we tested whether Brownian or Ornstein-Uhlenbeck models best fitted the data. Here, we used fitContinuous function from geiger package (Harmon, Weir, Brock, Glor, & Challenger, 2008) in an R statistical environment (R Core Team, 2017). We collected AIC (Akaike information criterion, Akaike, 1974) values for the regressions of each evolutionary model applied to each of the 9999 phylogenetic trees.

We then calculated  $\Delta AIC$ , pairwise, between the evolutionary models, for each tree. Based on the distribution of 9999  $\Delta AIC$  values (mean  $\Delta AIC$   $0.48 \pm 0.53$ ), Brownian and Ornstein-Uhlenbeck models were equally supported. Given that analyses based on small sample sizes tend to spuriously support the more complex Ornstein-Uhlenbeck model (Cooper, Thomas, Venditti, Meade, & Freckleton, 2016), we assumed that Brownian evolution model adequately suited our data and were thus able to use for model fitting MCMCglmm R package (Hadfield, 2010; Hadfield & Nakagawa, 2010).

We assessed the relationships among the circannual deviation, species-specific annual survival rate, and time spent in a constant photoregime (cycle 0 versus cycle 1), controlling for random effects of individuals, species, seasonal activities and studies. To do so we fitted Bayesian mixed models in the package MCMCglmm in R. To account for the uncertainty in phylogenetic relationships, we first randomly sampled 100 trees from the downloaded 9999 phylogenetic trees. Since MCMCglmm handles only one phylogenetic tree per model, we further incorporated the subset in 100 MCMCglmm models within the package mulTree

(Guillerme & Healy, 2017; Gutiérrez, Rakhimberdiev, Piersma, & Thieltges, 2017).

Technical information on the model parameters used and the replicable code are provided in Appendix S6 (<https://git.io/vAELZ>).

We used DIC (deviance information criterion by Spiegelhalter, Best, Carlin, & van der Linde (2002) for model selection. Following Zuur, Ieno, & Smith (2007), we started model selection from the most complete model:

$$Deviation_{BMR} \sim Survival_{BMR} + Cycle\_number + Survival_{BMR}:Cycle\_number$$

*random effects: LCS+study+species+species:BirdID + pedigree,*

where  $Deviation_{BMR}$  is the circannual deviation corrected for species-specific BMR,  $Survival_{BMR}$  is the effect of species-specific annual survival rate corrected for species-specific BMR,  $Cycle\_number$  distinguishes between the effects of transitional cycle and the cycle fully undergone under experimental photoregime,  $Survival_{BMR}:Cycle\_number$  is an interaction term.  $Species:BirdID$ ,  $LCS$ ,  $study$ ,  $species$  and  $pedigree$  define random effects for the: individual bird, life-cycle stage, study, from which the data were retrieved, bird species and phylogenetic variable, respectively. None of the random effects are nested within each other.

We calculated an analogue of Pagel's  $\lambda$ , or phylogenetic signal (the amount of variance explained by the phylogeny compared to the total amount of among species variance), of annual chronotype as  $var(phylogenetic)/(var(phylogenetic)+var(species))$ , phylogenetic variance contribution (the amount of variance explained by the phylogeny compared to the total variation in the data) following Bulla *et al.* (2016) as:

$$var(phylogenetic)/(var(phylogenetic)+var(BirdID)+var(LSC)+var(study)+var(species)+var(residual)),$$

with an intra-specific agreement repeatability as:

$$\text{var}(\text{species}) / (\text{var}(\text{species}) + \text{var}(\text{phylogenetic}) + \text{var}(\text{BirdID}) + \text{var}(\text{LSC}) + \text{var}(\text{study}) + \text{var}(\text{residual})),$$

and the adjusted repeatability as:  $\text{var}(\text{species}) / (\text{var}(\text{species}) + \text{var}(\text{residual}))$  following (Nakagawa & Schielzeth, 2010).

A small sample size may cause bias in a model accounting for phylogeny (Garamszegi & Møller, 2010). Thus, we provide code and an output of an MCMCglmm model of the same parameter structure, but not including the phylogenetic variable, in Appendices S6 and S7 (<https://git.io/vAELZ> and <https://git.io/vAuHK>).

## Results

Deviation of circannual periods in the species included in this study varied between 170 and 635 days, with a mean value of 371 days, and annual survival rates ranged from 0.28 to 0.90 with a mean = 0.62. Consistent with our prediction, the BMR-adjusted circannual periods were longer in species with higher BMR-adjusted annual survival rates. The correlation slope was steeper during the transition from the natural to an unvarying photoperiod (cycle 0: *mean slope estimate* = 215.1 days, 95% credible interval between 63.8 and 369.5 days) than the first cycle under such conditions (cycle 1: *mean slope estimate* = 133.4 days, 95% credible interval between -13.9 and 282.0 days; Fig. 1, Appendix S7 <https://git.io/vAuHK>). The difference in slopes between cycle 0 and cycle 1 was statistically significant, as indicated by  $\Delta\text{DIC} = 8.2$ , and, therefore, the full set of parameters was also the best one (Appendix S7).

The phylogenetic signal  $\lambda$  for circannual period length was 0.81 and the phylogenetic variance contribution was 0.43. The adjusted and agreement intra-specific repeatabilities of the trait were 0.95 and 0.10, respectively. The slope of correlation between annual survival rate and circannual period and its statistical significance were similar in the models that

included or did not account for phylogenetic effect (cycle 0: *mean slope estimate* = 200.2 days, 0.95% credible interval between 73.3 and 331.8 days; cycle 1: *mean slope estimate* = 120.3 days, 95% credible interval between - 0.04 and 247.2 days, Appendix S7).

## Discussion

It has long been known that free-running circannual periods in living organisms vary in length and rarely match the 365-day year (Gwinner, 1986). However, it was only recently proposed that the variation in circannual periods is adaptive; polymorphism in annual chronotypes facilitates a good match between the life histories and environments of organisms (Helm et al., 2017; Helm & Lincoln, 2017; Helm & Shavit, 2017). Here, we, for the first time, provide correlative evidence suggesting that annual chronotypes are the outcome of endogenous-clock properties having been adjusted to selective environments. We test the hypothesis that deviations of the circannual-clock cycle from the calendar year actually reflect solutions for the timing-mediated life-history trade-offs.. Consistent with our hypothesis, circannual periods were shorter in the species with lower probabilities of annual survival.

Our analysis was limited to migratory birds breeding in temperate and Arctic zones of the northern hemisphere. Selection pressures on the fitness components can be differently distributed along the time-axis, depending on migratory strategy or geographic and climatic factors for example (Jonzén, Hedenström, & Lundberg, 2007; Kristensen *et al.*, 2015). Thus, it is possible that the patterns reported here will not be upheld for resident bird populations, or for species breeding in the southern hemisphere or in the tropics (Newton & Brockie, 2008; Winkler, Ringelman, *et al.*, 2014; Young, 1994). Nevertheless, it seems unlikely that annual chronotypes will not work in concert with other life-history correlates, as long as there are seasonal selection pressures on fitness parameters.

## Variable circannual periods

In nature, variable environmental conditions require plasticity in life-history strategies (West-Eberhard, 2003). For this reason, the mechanisms coordinating organismal and environmental processes such as stress response or circadian chronotype may vary in the course of development, adjusting to environmental conditions experienced by individuals (Del Giudice, Ellis, & Shirtcliff, 2011; Roenneberg *et al.*, 2004). As a consequence, the coordinating mechanisms will also vary among individuals with different backgrounds or going through different ontogenetic and life-cycle stages. Our study, as well as many previous studies on circannual rhythms in birds (Helm, 2006), revealed high variation in circannual-period lengths among individuals of the same species, but also within each individual. For instance, in shorebirds, circannual periods of spring activities were closer to 365 days than the periods of activities expressed later in the season (Karagicheva, Rakhimberdiev, *et al.*, 2016; Piersma *et al.*, 2008). Karagicheva, Rakhimberdiev, *et al.* (2016) suggested that variations in the precision of expression of circannual cycles in different traits reflect selection pressures for precise timing changing with season.

Timing-dependent selection pressures will also vary with life style. For example, short circannual periods (and early chronotypes) should be useful for the opportunistically breeding red crossbill *Loxia curvirostra*, as they have to be alert to environmental cues almost year-round to opportunistically prepare for nesting as soon as pine cones become available as a food source (Hahn, 1998). In house sparrows *Passer domesticus* (not included in the analysis), the average circannual periods exerted in 12:12 LD are relatively short (10 months according to Wikelski *et al.*, 2008) for a species annual survival rate of 0.55 (Siriwardena *et al.*, 1999). Such a short circannual period is consistent with gonadal recrudescence starting in late autumn in house sparrows, perhaps an adaptation to high territorial competition in a sedentary bird (Hegner & Wingfield, 1986). In contrast, precise

Accepted Article

timing will be particularly important for long-distance migrants, and especially extreme “long-jump” migrants, of which Arctic-breeding shorebirds are a typical example (Åkesson *et al.*, 2017; Bazzi *et al.*, 2016; Conklin, Senner, Battley, & Piersma, 2017; Helm, Gwinner, & Trost, 2005; Piersma, 1987). For these species, environmental cues at departure from equatorial or temperate wintering sites are uninformative of the phenological conditions at remote Arctic breeding grounds (Piersma, *et al.*, 1990; Winkler, Jørgensen, *et al.*, 2014). To time their spring migration, these shorebirds largely rely on endogenous clocks, which have shown to be remarkably individually precise both in the wild as under unvarying photoperiodic conditions in captivity (Conklin, Battley, & Potter, 2013; Karagicheva, Rakhimberdiev, *et al.*, 2016).

The high variation in circannual deviations was well captured by the set of random variables included in our model. As a result, the unexplained (residual) variance was low, which caused the values of phylogenetic signal and intra-specific repeatability to be high when they were calculated accounting for the residual variance only, and contrastingly low when all random variance was taken into account. The difference suggests that circannual chronotype is a heritable, but highly variable, trait. However, we suggest caution when interpreting the phylogeny-related results in this paper, as our analysis was based on a small sample of 16 species from two orders (Garamszegi & Møller, 2010). The phylogenetic correlation was included in the model only to ensure that the hypothesised correlation was not confirmed spuriously due to bias from shared ancestry. Our key finding was that even when shared ancestry was accounted for, the correlation between length of circannual period and annual survival rate remained significant.

## Annual chronotype as a life-history covariate

Self-sustaining physiological oscillations of different periodicity in organisms, such as circadian, lunar, tidal or circannual rhythms, play an important role in coordinating rate processes (Hut, Paolucci, Dor, Kyriacou, & Daan, 2013; Neill, 2013; Numata & Helm, 2014; Riede *et al.*, 2017). While circannual oscillations are affected and probably even driven by internal metabolic processes (Wikelski *et al.*, 2008), they are also responsive to external information on environmental seasonal state received from the cue-response systems (Gwinner, 2003; Hahn & MacDougall-Shackleton, 2008). On this basis, we hypothesised that annual chronotypes of migratory birds should correlate with their other life-history traits, and specifically with species-specific annual survival probabilities. Despite vast variation in circannual periods for reasons of individuality, life-style, environment and ancestry (discussed above), we captured this relationship in the assembled data.

Our results extend the view that endogenous rhythms interact with external cues to ensure optimal adjustment of organisms to environments by modulation of the rate of entrainment of endogenous cycles to environmental oscillations (Floessner & Hut, 2017; Helm *et al.*, 2017; Helm & Lincoln, 2017). We projected circannual chronotypes on the slow-fast life-history axis and showed how the flexible system of endogenous oscillations helps birds to cope with temporal variation in environment in ways that are most appropriate for their life-history and life-table attributes. We propose that physiological circannual clocks, being responsive to the endogenous organismal and environmental processes at the same time, can be a physiological mediator coordinating energy in metabolic processes with survival and other organismal properties by adjusting them to external conditions through physiological and behavioural responses.



## Authors' contributions

J.K. assembled data and performed the statistical analyses. E.R. and T.P. participated in the concept development. E.R. and A.S. advised on the analyses. J.K. and T.P. wrote the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

## Acknowledgements

We are grateful to Robert Ricklefs and Roelof Hut for their constructive comments on earlier versions of the manuscript, to Martin Bulla, Jorge S. Gutiérrez and Pieternella Luttikhuisen for advice regarding phylogenetic analyses, to Jesse Conklin and Yvonne Verkuil for their help answering some questions from the reviewers, and to Dmitry Dorofeev for arranging access to the libraries in Russia. We thank Duncan Irschick and Charles Fox for their editorial work and and Brett K. Sandercock for his scrupulous edits and valuable comments. We thank the four unknown reviewers for improvement and thoughtful suggestions and questions. We deeply appreciated the expert opinions of Barbara Helm and William Schwartz, and thank them for their helpful remarks on the revised version of the manuscript. Our study was funded by the Spinoza Premium 2014 awarded to TP by the Netherlands Organisation for Scientific Research (NWO).

## Data Accessibility

Data, reproducible R-code and model results is also maintained on GitHub ([https://github.com/JKaragicheva/Circannual\\_cycles](https://github.com/JKaragicheva/Circannual_cycles)) and publicly archived at <https://doi.org/10.5281/zenodo.1250826>.

## References

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, **19**, 716–723. doi:10.1109/TAC.1974.1100705
- Åkesson, S., Ilieva, M., Karagicheva, J., Rakhimberdiev, E., Tomotani, B., & Helm, B. (2017). Timing avian long-distance migration: from internal clock mechanisms to global flights. *Philosophical Transactions of the Royal Society B*, **372**, 20160252. doi:10.1098/rstb.2016.0252
- Alonso, D., & Arizaga, J. (2013). The impact of vagrants on apparent survival estimation in a population of common crossbills (*Loxia curvirostra*). *Journal of Ornithology*, **154**, 209–217. doi:10.1007/s10336-012-0887-2
- Battley, P. F., Dekinga, A., Dietz, M. W., Piersma, T., Tang, S., & Hulsman, K. (2001). Basal metabolic rate declines during long-distance migratory flight in great knots. *Condor*, **103**, 838–845. doi:10.1650/0010-5422(2001)103[0838:BMRDDL]2.0.CO;2
- Bazzi, G., Cecere, J. G., Caprioli, M., Gatti, E., Gianfranceschi, L., Podofillini, S., ... Rubolini, D. (2016). Clock gene polymorphism, migratory behaviour and geographic distribution: a comparative study of trans-Saharan migratory birds. *Molecular Ecology*, **25**, 6077–6091. doi:10.1111/mec.13913
- Berthold, P. (1977). Über eine mögliche endogene Steuerung der Zugdisposition beim Fichtenkreuzschnabel *Loxia curvirostra*. *Journal of Ornithology*, **118**, 203–204. doi:10.1007/BF01648323
- Berthold, P. (1982). Endogene Grundlagen der Jahresperiodik von Standvögeln und wenig ausgeprägten Zugvögeln. *Journal of Ornithology*, **123**, 1–17. doi:10.1007/BF01644145

Berthold, P., Gwinner, E., & Klein, H. (1972). Circannuale Periodik bei Grasmücken. I.

Periodik des Körpergewichtes, der Mauser und der Nachtunruhe bei *Sylvia atricapilla* und *S. borin* unter verschiedenen konstanten Bedingungen. *J Ornithol*, **113**, 186–189.

Bielby, J., Mace, G. M., Bininda-Emonds, O. R. P., Cardillo, M., Gittleman, J. L., Jones, K. E., ... Purvis, A. (2007). The fast-slow continuum in mammalian life history: an empirical reevaluation. *American Naturalist*, **169**, 748–757. doi:10.1086/516847

Bradshaw, W. E., & Holzapfel, C. M. (2007). Evolution of animal photoperiodism. *Annual Review of Ecology, Evolution, and Systematics*, **38**, 1–25.  
doi:10.1146/annurev.ecolsys.37.091305.110115

Bulla, M., Valcu, M., Dokter, A. M., Dondua, A. G., Kosztolányi, A., Rutten, A. L., ... Kempenaers, B. (2016). Unexpected diversity in socially synchronized rhythms of shorebirds. *Nature*, **540**, 109–113. doi:10.1038/nature20563

Chandola-Saklani, A., Thapliyal, A., Negi, K., Diyundi, S. C., & Choudhary, B. (2004). Daily increments of light hours near vernal equinox synchronize circannual testicular cycle of tropical spotted munia. *Chronobiology International*, **21**, 553–569.

COMADRE Animal Matrix Database. (2015). (Version 1.0.0). Germany: Max Planck Institute for Demographic Research. Retrieved from [www.comadre-db.org](http://www.comadre-db.org) on 9/8/2016.

Conklin, J. R., Battley, P. F., & Potter, M. A. (2013). Absolute consistency: individual versus population variation in annual-cycle schedules of a long-distance migrant bird. *PLoS ONE*, **8**, e54535. doi:10.1371/journal.pone.0054535

Conklin, J. R., Senner, N. R., Battley, P. F., & Piersma, T. (2017). Extreme migration and the individual quality spectrum. *Journal of Avian Biology*, **48**, 19–36.  
doi:10.1111/jav.01316

Cooper, N., Thomas, G. H., Venditti, C., Meade, A., & Freckleton, R. P. (2016). A cautionary note on the use of Ornstein-Uhlenbeck models in macroevolutionary studies.

*Biological Journal of the Linnean Society*, **118**, 64–77. doi:10.1111/bij.12701

Dawson, A. (2008). Control of the annual cycle in birds: endocrine constraints and plasticity in response to ecological variability. *Philosophical Transactions of the Royal Society B*, **363**, 1621–1633. doi:10.1098/rstb.2007.0004

Dawson, A. (2015). Annual gonadal cycles in birds: modeling the effects of photoperiod on seasonal changes in GnRH-1 secretion. *Frontiers in Neuroendocrinology*, **37**, 52–64. doi:10.1016/j.yfrne.2014.08.004

Dawson, A., King, V. M., Bentley, G. E., & Ball, G. F. (2001). Photoperiodic control of seasonality in birds. *Journal of Biological Rhythms*, **16**, 365–380. doi:10.1177/074873001129002079

Del Giudice, M., Ellis, B. J., & Shirlcliff, E. A. (2011). The Adaptive Calibration Model of stress responsivity. *Neuroscience & Biobehavioral Reviews*, **35**, 1562–1592. doi:10.1016/j.neubiorev.2010.11.007

Desholm, M. (2009). Avian sensitivity to mortality: prioritising migratory bird species for assessment at proposed wind farms. *Journal of Environmental Management*, **90**, 2672–2679. doi:10.1016/j.jenvman.2009.02.005

Downing, P. A., Cornwallis, C. K., & Griffin, A. S. (2015). Sex, long life and the evolutionary transition to cooperative breeding in birds. *Proceedings of the Royal Society B*, **282**, 20151663. doi:10.1098/rspb.2015.1663

Drent, R., Both, C., Green, M., Madsen, J., & Piersma, T. (2003). Pay-offs and penalties of competing migratory schedules. *Oikos*, **103**, 274–292. doi:10.1034/j.1600-0706.2003.12274.x

Drent, R., & Daan, S. (1980). The prudent parent: energetic adjustments in avian breeding.

*Ardea*, **68**, 225–252.

Ekman, J., & Askenmo, C. (1986). Reproductive cost, age-specific survival and a comparison of the reproductive strategy in two European tits (genus *Parus*). *Evolution*, **40**, 159–168.

Floessner, T., & Hut, R. A. (2017). Basic principles underlying biological oscillations and their entrainment. In V. Kumar (Ed.), *Biological timekeeping: clocks, rhythms and behaviour* (pp. 47–58). New Delhi: Springer India. doi:10.1007/978-81-322-3688-7\_3

Freeman, S. N., Robinson, R. A., Clark, J. A., Griffin, B. M., & Adams, S. Y. (2007).

Changing demography and population decline in the common starling *Sturnus vulgaris*: a multisite approach to integrated population monitoring. *Ibis*, **149**, 587–596. doi:10.1111/j.1474-919X.2007.00684.x

Garamszegi, L. Z., & Møller, A. P. (2010). Effects of sample size and intraspecific variation in phylogenetic comparative studies: a meta-analytic review. *Biological Reviews*, **85**, 797–805. doi:10.1111/j.1469-185X.2010.00126.x

Glazier, D. S. (2015). Is metabolic rate a universal ‘pacemaker’ for biological processes?

*Biological Reviews*, **90**, 377–407. doi:10.1111/brv.12115

Guillerme, T., & Healy. (2017). *mulTree: performs MCMCglmm on multiple phylogenetic trees*. R package version 1.2.5.

Gutiérrez, J. S., Rakhimberdiev, E., Piersma, T., & Thieltges, D. W. (2017). Migration and parasitism: habitat use, not migration distance, influences helminth species richness in Charadriiform birds. *Journal of Biogeography*, **44**, 1137–1147. doi:10.1111/jbi.12956

Gwinner, E. (1968). Circannuale Periodik als Grundlage des jahreszeitlichen

Funktionswandels bei Zugvögeln: Untersuchungen am Fitis (*Phylloscopus trochilus*)

und am Waldlaubsänger (*P. sibilatrix*). *Journal of Ornithology*, **109**, 70–95.

doi:10.1007/BF01678108

Gwinner, E. (1971). A comparative study of circannual rhythms in warblers. In M. Menaker (Ed.), *Biochronometry* (pp. 405–427). Washington, D. C.: National Academy of Sciences.

Gwinner, E. (1981). Circannual rhythms: their dependence on the circadian system. In *Biological clocks in seasonal reproductive cycles* (pp. 153–169). Bristol: Wright and Sons.

Gwinner, E. (1986). *Circannual rhythms: endogenous annual clocks in the organization of seasonal processes*. Berlin: Springer-Verlag.

Gwinner, E. (2003). Circannual rhythms in birds. *Current Opinion in Neurobiology*, **13**, 770–778. doi:10.1016/j.conb.2003.10.010

Gwinner, E., & Schwabl-Benzinger, I. (1982). Adaptive temporal programming of molt and migratory disposition in two closely related long-distance migrants, the pied flycatcher (*Ficedula hypoleuca*) and the collared flycatcher (*Ficedula albicollis*). In F. Papi (Ed.), *Avian Navigation* (pp. 75–89). Springer Berlin Heidelberg.

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*, **33**, 1–22.

Hadfield, J. D., & Nakagawa, S. (2010). General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology*, **23**, 494–508. doi:10.1111/j.1420-9101.2009.01915.x

Hahn, T. P. (1998). Reproductive seasonality in an opportunistic breeder, the red crossbill, *Loxia curvirostra*. *Ecology*, **79**, 2365–2375. doi:10.1890/0012-9658(1998)079[2365:RSIAOB]2.0.CO;2

- Hahn, T. P., & MacDougall-Shackleton, S. A. (2008). Adaptive specialization, conditional plasticity and phylogenetic history in the reproductive cue response systems of birds. *Philosophical Transactions of the Royal Society B*, **363**, 267–286. doi:10.1098/rstb.2007.2139
- Harmon, L. J., Weir, J. T., Brock, C. D., Glor, R. E., & Challenger, W. (2008). GEIGER: investigating evolutionary radiations. *Bioinformatics*, **24**, 129–131. doi:10.1093/bioinformatics/btm538
- Harts, A. M. F., Kristensen, N. P., & Kokko, H. (2016). Predation can select for later and more synchronous arrival times in migrating species. *Oikos*, **125**, 1528–1538. doi:10.1111/oik.02973
- Harvey, P. H., & Pagel, M. D. (1991). *The comparative method in evolutionary biology*. Oxford ; New York: Oxford University Press.
- Hau, M., Dominoni, D., Casagrande, S., Buck, C. L., Wagner, G., Hazlerigg, D., ... Hut, R. A. (2017). Timing as a sexually selected trait: the right mate at the right moment. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **372**, 20160249. doi:10.1098/rstb.2016.0249
- Hegner, R. E., & Wingfield, J. C. (1986). Behavioral and endocrine correlates of multiple brooding in the semicolonial house sparrow *Passer domesticus* I. Males. *Hormones and Behavior*, **20**, 294–312. doi:10.1016/0018-506X(86)90039-5
- Helm, B. (2006). *Zugunruhe* of migratory and non-migratory birds in a circannual context. *Journal of Avian Biology*, **37**, 533–540. doi:10.1111/j.2006.0908-8857.03947.x
- Helm, B., Gwinner, E., & Trost, L. (2005). Flexible seasonal timing and migratory behavior: results from stonechat breeding programs. *Annals of the New York Academy of Sciences*, **1046**, 216–227. doi:10.1196/annals.1343.019

- Helm, B., & Lincoln, G. A. (2017). Circannual rhythms anticipate the Earth's annual periodicity. In V. Kumar (Ed.), *Biological timekeeping: clocks, rhythms and behaviour* (pp. 545–569). Springer India.
- Helm, B., Piersma, T., & van der Jeugd, H. (2006). Sociable schedules: interplay between avian seasonal and social behaviour. *Animal Behaviour*, **72**, 245–262.  
doi:10.1016/j.anbehav.2005.12.007
- Helm, B., Schwabl, I., & Gwinner, E. (2009). Circannual basis of geographically distinct bird schedules. *Journal of Experimental Biology*, **212**, 1259–1269. doi:10.1242/jeb.025411
- Helm, B., & Shavit, A. (2017). Dissecting and reconstructing time and space for replicable biological research. In *Stepping in the Same River Twice: Replication in Biological Research* (Ellison, A. M. and Shavit, A., pp. 233–249). New Haven: Yale University Press.
- Helm, B., Visser, M. E., Schwartz, W., Kronfeld-Schor, N., Gerkema, M., Piersma, T., & Bloch, G. (2017). Two sides of a coin: ecological and chronobiological perspectives of timing in the wild. *Philosophical Transactions of the Royal Society B*, **372**, 20160246. doi:10.1098/rstb.2016.0246
- Holberton, R. L., & Able, K. P. (1992). Persistence of circannual cycles in a migratory bird held in constant dim light. *Journal of Comparative Physiology A*, **171**, 477–481.
- Huelsenbeck, J. P. (2000). Accommodating phylogenetic uncertainty in evolutionary studies. *Science*, **288**, 2349–2350. doi:10.1126/science.288.5475.2349
- Hut, R. A., Paolucci, S., Dor, R., Kyriacou, C. P., & Daan, S. (2013). Latitudinal clines: an evolutionary view on biological rhythms. *Proceedings of the Royal Society B*, **280**, 20130433–20130433. doi:10.1098/rspb.2013.0433
- Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, **491**, 444–448. doi:10.1038/nature11631



Johnston, A., Robinson, R. A., Gargallo, G., Julliard, R., van der Jeugd, H., & Baillie, S. R.

(2016). Survival of Afro-Palaeartic passerine migrants in Western Europe and the impacts of seasonal weather variables. *Ibis*, **158**, 465–480. doi:10.1111/ibi.12366

Jonzén, N., Hedenström, A., & Lundberg, P. (2007). Climate change and the optimal arrival of migratory birds. *Proceedings of the Royal Society B*, **274**, 269–274. doi:10.1098/rspb.2006.3719

Karagicheva, J., Liebers, M., Rakhimberdiev, E., Hallinger, K. K., Saveliev, A., & Winkler, D. W. (2016). Differences in size between first and replacement clutches match the seasonal decline in single clutches in tree swallows *Tachycineta bicolor*. *Ibis*, **158**, 607–613. doi:10.1111/ibi.12368

Karagicheva, J., Rakhimberdiev, E., Dekinga, A., Brugge, M., Koolhaas, A., ten Horn, J., & Piersma, T. (2016). Seasonal time keeping in a long-distance migrating shorebird. *Journal of Biological Rhythms*, **31**, 509–521. doi:10.1177/0748730416655929

Kokko, H. (1999). Competition for early arrival in migratory birds. *Journal of Animal Ecology*, **68**, 940–950. doi:10.1046/j.1365-2656.1999.00343.x

Kristensen, N. P., Johansson, J., Ripa, J., & Jonzén, N. (2015). Phenology of two interdependent traits in migratory birds in response to climate change. *Proceedings of the Royal Society B*, **282**, 20150288–20150288. doi:10.1098/rspb.2015.0288

Lok, T., Veldhoen, L., Overdijk, O., Tinbergen, J. M., & Piersma, T. (2017). An age-dependent fitness cost of migration? Old trans-Saharan migrating spoonbills breed later than those staying in Europe, and late breeders have lower recruitment. *Journal of Animal Ecology*, **86**, 998–1009. doi:10.1111/1365-2656.12706

McGraw, J. B., & Caswell, H. (1996). Estimation of individual fitness from life-history data. *American Naturalist*, **1**, 47–64.

- McKechnie, A. E. (2008). Phenotypic flexibility in basal metabolic rate and the changing view of avian physiological diversity: a review. *Journal of Comparative Physiology B*, **178**, 235–247. doi:10.1007/s00360-007-0218-8
- McNamara, J. M., Welham, R. K., Houston, A. I., Daan, S., & Tinbergen, J. M. (2004). The effects of background mortality on optimal reproduction in a seasonal environment. *Theoretical Population Biology*, **65**, 361–372. doi:10.1016/j.tpb.2003.10.006
- Møller, A. P. (1994). Phenotype-dependent arrival time and its consequences in a migratory bird. *Behavioral Ecology and Sociobiology*, **35**, 115–122.
- Møller, A. P. (2008). Relative longevity and field metabolic rate in birds. *Journal of Evolutionary Biology*, **21**, 1379–1386. doi:10.1111/j.1420-9101.2008.01556.x
- Møller, A. P. (2009). Basal metabolic rate and risk-taking behaviour in birds: BMR and anti-predator behaviour. *Journal of Evolutionary Biology*, **22**, 2420–2429. doi:10.1111/j.1420-9101.2009.01850.x
- Moreno, J., Gustafsson, L., Carlson, A., & Pärt, T. (1991). The cost of incubation in relation to clutch-size in the collared flycatcher *Ficedula albicollis*. *Ibis*, **133**, 186–193. doi:10.1111/j.1474-919X.1991.tb04830.x
- Morrison, C. A., Robinson, R. A., Butler, S. J., Clark, J. A., & Gill, J. A. (2016). Demographic drivers of decline and recovery in an Afro-Palaeartic migratory bird population. *Proceedings of the Royal Society B: Biological Sciences*, **283**, 20161387. doi:10.1098/rspb.2016.1387
- Mueller, M., Spaar, R., Schifferli, L., & Jenni, L. (2005). Effects of changes in farming of subalpine meadows on a grassland bird, the whinchat (*Saxicola rubetra*). *Journal of Ornithology*, **146**, 14–23. doi:10.1007/s10336-004-0059-0

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists. *Biological Reviews*, **85**, 935–956. doi:10.1111/j.1469-185X.2010.00141.x

Neill, D. (2013). Life's timekeeper. *Ageing Research Reviews*, **12**, 567–578. doi:10.1016/j.arr.2013.01.004

Newton, I. (2008). *The migration ecology of birds*. Amsterdam: Elsevier/Acad. Press.

Numata, H., & Helm, B. (Eds.). (2014). *Annual, lunar, and tidal clocks: patterns and mechanisms of nature's enigmatic rhythms*. Tokyo ; New York: Springer.

Öberg, M., Pärt, T., Arlt, D., Laugen, A. T., & Low, M. (2014). Decomposing the seasonal fitness decline. *Oecologia*, **174**, 139–150. doi:10.1007/s00442-013-2763-z

Orzack, S. H., & Tuljapurkar, S. (2001). Reproductive effort in variable environments, or environmental variation is for the birds. *Ecology*, **82**, 2659–2665. doi:10.1890/0012-9658(2001)082[2659:REIVEO]2.0.CO;2

Piersma, T. (1987). Hop, skip, or jump? Constraints on migration of arctic waders by feeding, fattening, and flight speed. *Limosa*, **60**, 185–194.

Piersma, T., Klaassen, M., Bruggemann, J.H., Blomert, A.-M., Gueye, A., Ntiamoa-Baidu, Y., & van Brederode, N.E. (1990). Seasonal timing of the spring departure of waders from the Banc d'Arguin, Mauritania. *Ardea*, **78**, 123-134.

Piersma, T., Brugge, M., Spaans, B., & Battley, P. F. (2008). Endogenous circannual rhythmicity in body mass, molt, and plumage of great knots (*Calidris tenuirostris*). *Auk*, **125**, 140–148. doi:10.1525/auk.2008.125.1.140

Piersma, T., Cadée, N., & Daan, S. (1995). Seasonality in basal metabolic rate and thermal conductance in a long distance migrant shorebird, the knot (*Calidris canutus*). *Journal of Comparative Physiology*, **165**, 37–45.

Piersma, T., Lok, T., Chen, Y., Hassell, C. J., Yang, H.-Y., Boyle, A., ... Ma, Z. (2016).

Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *Journal of Applied Ecology*, **53**, 479–490. doi:10.1111/1365-2664.12582

Pohl, H. (1971). Circannuale Periodik beim Bergfinken. *Naturwissenschaften*, 58(11), 572–573.

R Core Team. (2017). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>

Rakhimberdiev, E., van den Hout, P. J., Brugge, M., Spaans, B., & Piersma, T. (2015). Seasonal mortality and sequential density dependence in a migratory bird. *Journal of Avian Biology*, **46**, 332–341. doi:10.1111/jav.00701

Rani, S., & Kumar, V. (2013). Avian circannual systems: Persistence and sex differences. *General and Comparative Endocrinology*, **190**, 61–67. doi:10.1016/j.ygcen.2013.04.013

Reale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P. O. (2010). Personality and the emergence of the pace-of-life syndrome concept at the population level. *Philosophical Transactions of the Royal Society B*, **365**, 4051–4063. doi:10.1098/rstb.2010.0208

Ricklefs, R. E., & Wikelski, M. (2002). The physiology/life-history nexus. *Trends in Ecology & Evolution*, **17**, 462–468. doi:10.1016/S0169-5347(02)02578-8

Riede, S. J., van der Vinne, V., & Hut, R. A. (2017). The flexible clock: predictive and reactive homeostasis, energy balance and the circadian regulation of sleep–wake timing. *Journal of Experimental Biology*, **220**, 738–749. doi:10.1242/jeb.130757

Roenneberg, T., Daan, S., & Mrosovsky, M. (2003). The art of entrainment. *Journal of Biological Rhythms*, **18**, 183–194. doi:10.1177/0748730403018003001

- Accepted Article
- Roenneberg, T., Kuehnle, T., Pramstaller, P. P., Ricken, J., Havel, M., Guth, A., & Mellow, M. (2004). A marker for the end of adolescence. *Current Biology*, **14**, R1038–R1039. doi:10.1016/j.cub.2004.11.039
- Sæther, B.-E., & Bakke, Ø. (2000). Avian life history variation and contribution of demographic traits to the population growth rate. *Ecology*, **81**, 642–653. doi:10.1890/0012-9658(2000)081[0642:ALHVAC]2.0.CO;2
- Sanz, J. J. (2001). Latitudinal variation in female local return rate in the philopatric pied flycatcher (*Ficedula hypoleuca*). *Auk*, **118**, 539. doi:10.1642/0004-8038(2001)118[0539:LVIFLR]2.0.CO;2
- Senar, J. ., Borrás, A., Cabrera, T., & Cabrera, J. (1993). Testing for the relationship between coniferous crop stability and common crossbill residence. *Journal of Field Ornithology*, **64**, 464–469.
- Silverin, B., Arvidsson, B., & Wingfield, J. (1997). The adrenocortical responses to stress in breeding willow warblers *Phylloscopus trochilus* in Sweden: effects of latitude and gender. *Functional Ecology*, **11**, 376–384. doi:10.1046/j.1365-2435.1997.00097.x
- Siriwardena, G. M., Baillie, S. R., & Wilson, J. D. (1998). Variation in the survival rates of some British passerines with respect to their population trends on farmland. *Bird Study*, **45**, 276–292. doi:10.1080/00063659809461099
- Siriwardena, G. M., Baillie, S. R., & Wilson, J. D. (1999). Temporal variation in the annual survival rates of six granivorous birds with contrasting population trends. *Ibis*, **141**, 621–636. doi:10.1111/j.1474-919X.1999.tb07370.x
- Speakman, J. R. (2005). Correlations between physiology and lifespan - two widely ignored problems with comparative studies. *Aging Cell*, **4**, 167–175. doi:10.1111/j.1474-9726.2005.00162.x

- Spiegelhalter, D. J., Best, N. G., Carlin, B. P., & van der Linde, A. (2002). Bayesian measures of model complexity and fit. *Journal of the Royal Statistical Society B*, **64**, 583–639. doi:10.1111/1467-9868.00353
- van Wijk, R. E., Schaub, M., & Bauer, S. (2017). Dependencies in the timing of activities weaken over the annual cycle in a long-distance migratory bird. *Behavioral Ecology and Sociobiology*, **71**, 73. doi:10.1007/s00265-017-2305-5
- Versteegh, M. A., Schwabl, I., Jaquier, S., & Tieleman, B. I. (2012). Do immunological, endocrine and metabolic traits fall on a single pace-of-life axis? Covariation and constraints among physiological systems. *Journal of Evolutionary Biology*, **25**, 1864–1876. doi:10.1111/j.1420-9101.2012.02574.x
- Ward, S. (2004). Singing is not energetically demanding for pied flycatchers, *Ficedula hypoleuca*. *Behavioral Ecology*, **15**, 477–484. doi:10.1093/beheco/arh038
- Weiser, E. L., Brown, S. C., Lanctot, R. B., Gates, H. R., Abraham, K. F., Bentzen, R. L., ... Sandercock, B. K. (2017). Life-history tradeoffs revealed by seasonal declines in reproductive traits of Arctic-breeding shorebirds. *Journal of Avian Biology*. doi:10.1111/jav.01531
- West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. New York: Oxford University Press.
- Wikelski, M., Martin, L. B., Scheuerlein, A., Robinson, M. T., Robinson, N. D., Helm, B., ... Gwinner, E. (2008). Avian circannual clocks: adaptive significance and possible involvement of energy turnover in their proximate control. *Philosophical Transactions of the Royal Society B*, **363**, 411–423. doi:10.1098/rstb.2007.2147
- Williams, T. D. (2012). *Physiological adaptations for breeding in birds*. Princeton, N.J.: Princeton University Press.

Winkler, D. W., Jørgensen, C., Both, C., Houston, A. I., McNamara, J. M., Levey, D. J., ...

Piersma, T. (2014). Cues, strategies, and outcomes: how migrating vertebrates track environmental change. *Movement Ecology*, **2**, 10. doi:10.1186/2051-3933-2-10

Winkler, D. W., Ringelman, K. M., Dunn, P. O., Whittingham, L., Hussell, D. J. T., Clark, R.

G., ... Ardia, D. R. (2014). Latitudinal variation in clutch size-lay date regressions in *Tachycineta* swallows: effects of food supply or demography? *Ecography*, **37**, 670–678. doi:10.1111/j.1600-0587.2013.00458.x

Young, B. E. (1994). Geographic and seasonal patterns of clutch-size variation in house wrens. *The Auk*, **111**, 545–555.

Zuur, A. F., Ieno, E. N., & Smith, G. M. (2007). *Analysing ecological data*. New York; London: Springer. Retrieved from <http://public.eblib.com/choice/publicfullrecord.aspx?p=371384>

Дольник, В. (1975). Окологодовая цикличность миграционного отложения жира, половой активности и линьки при постоянных фотопериодах у зяблика (*Fringilla coelebs*). *Журн. Общ. Биол.*, (19746), 543–552.

#### SUPPORTING INFORMATION

Table S1 Circannual cycle lengths

Table S2 Species-specific annual survival rates

Table S3 Species-specific basal metabolic rates

Table S4 Species-specific body mass

Figure S5 Correlation BMR body mass

Appendix S6 R code

Table S7 MCMCglmm models output

Please note: Wiley Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.



Table 1. The circannual deviations from 365 days (mean and SD of the raw data points that were used in the analysis) and details on the experimental setups; species-specific annual survival probability and BMR.

Species	Number of individuals used in the experiment	Circannual deviation, days (mean $\pm$ SD)	Cycles since transition to unvarying photoregime	Photoperiod	Annual survival rate (median)	BMR, W	Body mass, g
<i>Ficedula albicollis</i>	10 males	28.6 $\pm$ 22.2 <sup>1</sup>	0,1	13:11 LD	0.45 <sup>14</sup>	0.276 <sup>34</sup>	14.2 <sup>14</sup>
<i>Ficedula hypoleuca</i>	7 males	36.3 $\pm$ 25.8 <sup>1</sup>	0,1	12:12 LD	0.48 <sup>15</sup>	0.230 <sup>35</sup>	11.6 <sup>14</sup>
<i>Fringilla coelebs</i>	15 males	53.5 $\pm$ 37.4 <sup>2</sup>	1	12:12 LD	0.6 <sup>16, 17</sup>	0.373 <sup>30</sup>	24.2 <sup>30</sup>
<i>Fringilla montifringilla</i>	2 males	90.0 $\pm$ 34.6 <sup>3</sup>	1	10:14 LD	0.57 <sup>18</sup>	0.516 <sup>31</sup>	24.1 <sup>31</sup>
<i>Junco hyemalis</i>	7 males	100.4 $\pm$ 83.6 <sup>4</sup>	1	24 DD	0.49 <sup>14</sup>	0.192 <sup>31</sup>	19.6 <sup>14</sup>
<i>Loxia curvirostra</i>	3 males	- 71.6 $\pm$ 54.2 <sup>5, 6</sup>	1	12:12 LD	0.45 <sup>19,20</sup>	0.600 <sup>30</sup>	40.6 <sup>30</sup>
<i>Phylloscopus trochilus</i>	6 unknown	- 22.3 $\pm$ 25.2 <sup>7, 8</sup>	0,1	12:12 LD	0.34 <sup>16,21,22,23</sup>	0.208 <sup>30</sup>	9.4 <sup>30</sup>
<i>Phylloscopus collybita</i>	7 unknown	- 10.0 $\pm$ 18.0 <sup>8</sup>	0	12:12 LD	0.28 <sup>16,22</sup>	0.165 <sup>30</sup>	7.7 <sup>30</sup>
<i>Parus cristatus</i>	8 unknown	- 46.3 $\pm$ 17.8 <sup>6</sup>	0,1	10:14 LD	0.6 <sup>14,24</sup>	0.470 <sup>14</sup>	11.2 <sup>14</sup>
<i>Passer montanus</i>	3 females	- 87.0 $\pm$ 66.6 <sup>6</sup>	0,1	10:14 LD and 14:10 LD	0.44 <sup>25</sup>	0.405 <sup>30</sup>	21.7 <sup>30</sup>
<i>Saxicola torquata rubicola</i>	9 males	- 30.0 $\pm$ 32.7 <sup>9</sup>	1	12.25:11.75	0.48 <sup>26</sup>	0.256 <sup>30</sup>	14.9 <sup>30</sup>

				LD			
<i>Sturnus vulgaris</i>	18 males	$33.7 \pm 74.6^{10}$	1	12:12 LD	0.72 <sup>14,16,27</sup>	$0.877^{30}$	$82.3^{14}$
<i>Sylvia borin</i>	4 males	$-25.7 \pm 61.5^{10,11}$	1	12:12 LD	$0.46^{16,22}$	$0.416^{30}$	$19.1^{30}$
<i>Sylvia atricapilla</i>	4 males	$-51.7 \pm 27.2^{11}$	1	12:12 LD	$0.41^{16,22}$	$0.413^{30}$	$18.9^{30}$
<i>Calidris canutus</i>	16 unknown	$6.8 \pm 37.3^{12}$	0,1	12:12 LD	$0.81^{28}$	$0.880^{32}$	$130.0^{32}$
<i>Calidris tenuirostris</i>	4 unknown	$8.7 \pm 45.6^{13}$	0,1	12:12 LD	$0.9^{29}$	$1.255^{33}$	$149.3^{33}$

1. Gwinner & Schwabl-Benzinger, 1982; 2. Дольник, 1975; 3. Pohl, 1971; 4. Holberton & Able, 1992; 5. Berthold, 1977; 6. Berthold, 1982; 7. Gwinner, 1968; 8. Gwinner, 1971; 9. Helm et al., 2009; 10. Gwinner, 1981; 11. Berthold, Gwinner, & Klein, 1972; 12. Karagicheva, Rakhimberdiev, et al., 2016; 13. Piersma, Brugge, Spaans, & Battley, 2008; 14. Møller, 2008; 15. Sanz, 2001; 16. Siriwardena, Baillie, & Wilson, 1998; 17. Siriwardena, Baillie, & Wilson, 1999; 18. Desholm, 2009; 19. Senar, Borrás, Cabrera, & Cabrera, 1993; 20. Alonso & Arizaga, 2013; 21. Silverin, Arvidsson, & Wingfield, 1997; 22. Johnston et al., 2016; 23. Morrison, Robinson, Butler, Clark, & Gill, 2016; 24. Ekman & Askenmo, 1986; 25. Downing, Cornwallis, & Griffin, 2015; 26. Mueller, Spaar, Schifferli, & Jenni, 2005; 27. Freeman, Robinson, Clark, Griffin, & Adams, 2007; 28. Rakhimberdiev, van den Hout, Brugge, Spaans, & Piersma, 2015; 29. Piersma et al., 2016; 30. Møller, 2009; 31. McKechnie, 2008; 32. Piersma, Cadée, & Daan, 1995; 33. Battley et al., 2001; 34. Moreno, Gustafsson, Carlson, & Pärt, 1991; 35. Ward, 2004.

